Page 3 of 13

Response to Restriction Requirement and Species Election

Applicant(s): BENZ et al. Scrial No.: 10/663,926 Confirmation No.: 2299 Filed: September 16, 2003

For: COMPOUNDS CONTAINING QUATERNARY CARBONS AND SILICON-CONTAINING GROUPS.

MEDICAL DEVICES, AND METHODS

## CLAIMS READABLE ON THE ELECTED SPECIES

1. A medical device comprising a polymer comprising a group of the formula:

$$-[-(R^1)_n-(-Z-(R^2)_n-)_n-(-Si(R)_2-V_r-)_s-]_{q^2}$$

wherein:

n = 0 or 1; m = 0 or 1; p = 1-100,000; r = 0-100,000; s = 1-100,000; q = 1-100,000;

R<sup>1</sup> and R<sup>2</sup> are each independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms;

Z is  $-C(R^3)_2$ - wherein each  $R^3$  is independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms, wherein the two  $R^3$  groups within  $-C(R^3)_2$ - can be optionally joined to form a ring;

each R is independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms; and

V is 
$$-O-Si(R)_2$$
- or  $R^1$ .

- 2. The medical device of claim 1 wherein p = 1-5000.
- 3. The medical device of claim 2 wherein p = 2-12.
- 4. The medical device of claim 1 wherein  $\mathbb{R}^1$  and  $\mathbb{R}^2$  are each independently a straight chain alkylene group, an arylene group, or combinations thereof.

Page 4 of 13

Response to Restriction Requirement and Species Election

Applicant(s): BENZ et al. Serial No.: 10/663,926 Confirmation No.: 2299 Filed: September 16, 2003

For: COMPOUNDS CONTAINING QUATERNARY CARBONS AND SILICON-CONTAINING GROUPS,

MEDICAL DEVICES, AND METHODS

5. The medical device of claim 4 wherein R<sup>1</sup> and R<sup>2</sup> are each independently a straight chain alkylene group.

- 6. The medical device of claim 1 wherein R<sup>1</sup> and R<sup>2</sup> are each independently groups containing up to 100 carbon atoms.
- 7. The medical device of claim 6 wherein R<sup>1</sup> and R<sup>2</sup> are each independently groups containing up to 20 carbon atoms.
- 8. The medical device of claim 7 wherein R<sup>1</sup> and R<sup>2</sup> are each independently groups containing 2 to 20 carbon atoms.
- 9. The medical device of claim 1 wherein each R<sup>3</sup> is independently a straight chain alkyl group, an aryl group, or combinations thereof, optionally including heteroatoms.
- 10. The medical device of claim 9 wherein each R<sup>3</sup> is independently a straight chain alkyl group, optionally including heteroatoms.
- 11. The medical device of claim 10 wherein each R<sup>3</sup> is independently a straight chain alkyl group containing 1 to 20 carbon atoms.
- 12. The medical device of claim 1 wherein the polymer further comprises a urethane group, a urea group, or combinations thereof.
- 13. The medical device of claim 12 wherein the polymer comprises a segmented polyurethane.
- 14. The medical device of claim 1 wherein the polymer is a biomaterial.

Page 5 of 13

Applicant(s): BENZ et al. Serial No.: 10/663,926 Confirmation No.: 2299 Filed: September 16, 2003

For: COMPOUNDS CONTAINING QUATERNARY CARBONS AND SILICON-CONTAINING GROUPS,

MEDICAL DEVICES. AND METHODS

- 15. The medical device of claim 14 wherein the polymer is substantially free of ether, ester, and carbonate linkages.
- 16. The medical device of claim 1 wherein the polymer is linear, branched, or crosslinked.
- 17. A medical device comprising a polymer prepared from a compound of the formula:

$$Y-[-(R^1)_n-(-Z-(R^2)_m-)_n-(-Si(R)_2-V_{r-1})_s-]_n-R^5-Y$$

wherein:

each Y is independently OH or NR<sup>4</sup>H;

n = 0 or 1;

m = 0 or 1;

p = 1-100,000;

r = 0-100,000;

s = 1-100,000;

q = 1-100,000;

R<sup>1</sup>, R<sup>2</sup>, and R<sup>5</sup> are each independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms;

Z is  $-C(R^3)_2$ - wherein each  $R^3$  is independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms, wherein the two  $R^3$  groups within  $-C(R^3)_2$ - can be optionally joined to form a ring;

each R is independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms;

each R<sup>4</sup> is independently H or a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof; and

V is 
$$-O-Si(R)_2$$
- or  $R'$ .

- 18. The medical device of claim 17 wherein p = 1-100.
- 19. The medical device of claim 18 wherein p = 2-12.

Page 6 of 13

Response to Restriction Requirement and Species Election

Applicant(s): BENZ et al. Serial No.: 10/663,926 Confirmation No.: 2299 Filed: September 16, 2003

For: COMPOUNDS CONTAINING QUATERNARY CARBONS AND SILICON-CONTAINING GROUPS,

MEDICAL DEVICES, AND METHODS

- 20. The medical device of claim 17 wherein the number average molecular weight of the compound of the formula  $Y-[-(R^1)_n-(-Z-(R^2)_m-)_p-(-Si(R)_z-V_r-)_x-]_q-R^5-Y$  is no greater than about 100,000 grams/mole.
- 21. The medical device of claim 20 wherein the number average molecular weight of the compound of the formula  $Y-[-(R^1)_n-(-Z-(R^2)_m-)_p-(-Si(R)_2-V_r-)_s-]_q-R^5-Y$  is about 1000 grams/mole to about 1500 grams/mole.
- 22. The medical device of claim 17 wherein R<sup>1</sup> and R<sup>2</sup> are each independently a straight chain alkylene group, an arylene group, or combinations thereof.
- 23. The medical device of claim 22 wherein R<sup>1</sup> and R<sup>2</sup> are each independently a straight chain alkylene group.
- 24. The medical device of claim 17 wherein R<sup>1</sup> and R<sup>2</sup> are each independently groups containing up to 100 carbon atoms.
- 25. The medical device of claim 24 wherein R<sup>1</sup> and R<sup>2</sup> are each independently groups containing up to 20 carbon atoms.
- 26. The medical device of claim 25 wherein R<sup>1</sup> and R<sup>2</sup> are each independently groups containing 2 to 20 carbon atoms.
- 27. The medical device of claim 17 wherein each R<sup>2</sup> includes at least two carbon atoms.
- 28. The medical device of claim 17 wherein each R<sup>3</sup> is independently a straight chain alkyl group, an aryl group, or combinations thereof, optionally including heteroatoms.
- 29. The medical device of claim 28 wherein each R<sup>3</sup> is independently a straight chain alkyl group, optionally including heteroatoms.

Page 7 of 13

Response to Restriction Requirement and Species Election

Applicant(s): BENZ et al. Serial No.: 10/663,926 Confirmation No.: 2299 Filed: September 16, 2003

For: COMPOUNDS CONTAINING QUATERNARY CARBONS AND SILICON-CONTAINING GROUPS,

MEDICAL DEVICES, AND METHODS

- 30. The medical device of claim 29 wherein each R<sup>3</sup> is independently a straight chain alkyl group containing 1 to 20 carbon atoms.
- 31. The medical device of claim 17 wherein the polymer further comprises a urethane group, a urea group, or combinations thereof.
- 32. The medical device of claim 31 wherein the polymer comprises a segmented polymethane.
- 33. The medical device of claim 17 wherein the polymer is a biomaterial.
- 34. The medical device of claim 33 wherein the polymer is substantially free of ether, ester, and carbonate linkages.
- 35. The medical device of claim 17 wherein each Y is OH.
- 36. The medical device of claim 17 wherein each R<sup>4</sup> is independently H or a straight chain alkyl group.
- 37. The medical device of claim 36 wherein each R<sup>4</sup> is independently a straight chain alkyl group containing 1 to 20 carbon atoms.
- 38. The medical device of claim 36 wherein each R4 is H.
- 39. The medical device of claim 17 wherein the polymer is linear, branched, or crosslinked.
- 40. A polymer comprising a group of the formula:

$$-[-(R^1)_n-(-Z-(R^2)_m-)_p-(-Si(R)_2-V_r-)_s-]_q-$$

Page 8 of 13

Applicant(s): BENZ et al. Serial No.: 10/663,926 Confirmation No.: 2299 Filed: September 16, 2003

For: COMPOUNDS CONTAINING QUATERNARY CARBONS AND SILICON-CONTAINING GROUPS,

MEDICAL DEVICES, AND METHODS

## wherein:

n = 0 or 1; m = 0 or 1; p = 1-100,000; r = 0-100,000; s = 1-100,000; q = 1-100,000;

R<sup>1</sup> and R<sup>2</sup> are each independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms;

Z is  $-C(R^3)_{2^-}$  wherein each  $R^3$  is independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms, wherein the two  $R^3$  groups within  $-C(R^3)_{2^-}$  can be optionally joined to form a ring;

each R is independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms; and

V is 
$$-O-Si(R)_{2}$$
 or  $R^{1}$ .

- 41. The polymer of claim 40 wherein p = 1-5000.
- 42. The polymer of claim 40 wherein p = 2-12.
- 43. The polymer of claim 40 wherein  $R^1$  and  $R^2$  are each independently a straight chain alkylene group, an arylene group, or combinations thereof.
- 44. The polymer of claim 43 wherein R<sup>1</sup> and R<sup>2</sup> are each independently a straight chain alkylene group.
- 45. The polymer of claim 40 wherein R<sup>1</sup> and R<sup>2</sup> are each independently groups containing 2 to 20 carbon atoms.

Page 9 of 13

Response to Restriction Requirement and Species Election

Applicant(s): BENZ et al. Serial No.: 10/663,926 Confirmation No.: 2299 Filed: September 16, 2003

For: COMPOUNDS CONTAINING QUATERNARY CARBONS AND SILICON-CONTAINING GROUPS,

MEDICAL DEVICES, AND METHODS

- 46. The polymer of claim 40 wherein each R<sup>3</sup> is independently a straight chain alkyl group, an aryl group, or combinations thereof, optionally including heteroatoms.
- 47. The polymer of claim 46 wherein each R<sup>3</sup> is independently a straight chain alkyl group, optionally including heteroatoms.
- 48. The polymer of claim 47 wherein each R<sup>3</sup> is independently a straight chain alkyl group containing 1 to 20 carbon atoms.
- 49. The polymer of claim 40 which is linear, branched, or crosslinked.
- 50. A polymer comprising a urethane group, a urea group, or combinations thereof, and a group of the formula:

$$-[-(R^1)_0-(-Z-(R^2)_0)-(-Si(R)_2-V_1-)_5-]_0$$

wherein:

n = 0 or 1; m = 0 or 1; p = 1-100,000; r = 0-100,000; s = 1-100,000; q = 1-100,000;

R<sup>1</sup> and R<sup>2</sup> are each independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms;

Z is  $-C(R^3)_2$ - wherein each  $R^3$  is independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms, wherein the two  $R^3$  groups within  $-C(R^3)_2$ - can be optionally joined to form a ring;

Page 10 of 13

Response to Restriction Requirement and Species Election

Applicant(s): BENZ et al. Serial No.: 10/663,926 Confirmation No.: 2299 Filed: September 16, 2003

For: COMPOUNDS CONTAINING QUATERNARY CARBONS AND SILICON-CONTAINING GROUPS,

MEDICAL DEVICES, AND METHODS

each R is independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms; and V is -O-Si(R)<sub>2</sub>- or R<sup>1</sup>.

- 51. The polymer of claim 50 wherein p = 1-100.
- 52. The polymer of claim 51 wherein p = 2-12.
- 53. The polymer of claim 50 which is a segmented polyurethane.
- 54. The polymer of claim 50 which is a biomaterial.
- 55. The polymer of claim 54 which is substantially free of ether, ester, and carbonate linkages.
- 56. The polymer of claim 50 which is linear, branched, or crosslinked.
- 57. A polymer prepared from a compound of the formula:

$$Y-[-(R^1)_n-(-Z-(R^2)_m-)_n-(-Si(R)_2-V_r-)_s-]_n-R^5-Y$$

wherein:

each Y is independently OH or NR⁴H;

n = 0 or 1;

m = 0 or 1;

p = 1-100,000;

r = 0-100,000;

s = 1-100,000;

q = 1-100,000;

Page 11 of 13

Response to Restriction Requirement and Species Election

Applicant(s): BENZ et al. Serial No.: 10/663.926 Confirmation No.: 2299

Confirmation No.: 2299 Filed: September 16, 2003

For: COMPOUNDS CONTAINING QUATERNARY CARBONS AND SILICON-CONTAINING GROUPS,

MEDICAL DEVICES, AND METHODS

R<sup>1</sup>, R<sup>2</sup>, and R<sup>5</sup> are each independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms;

Z is  $-C(R^3)_2$ - wherein each  $R^3$  is independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms, wherein the two  $R^3$  groups within  $-C(R^3)_2$ - can be optionally joined to form a ring;

each R is independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms;

each R<sup>4</sup> is independently H or a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof; and

V is 
$$-O-Si(R)_2$$
- or  $R^1$ .

- 58. The polymer of claim 57 wherein p = 1-100.
- 59. The polymer of claim 58 wherein p = 2-12.
- 60. The polymer of claim 57 wherein the number average molecular weight of the compound of the formula  $Y-[-(R^1)_n-(-Z-(R^2)_m-)_p-(-Si(R)_2-V_r-)_s-]_q-R^5-Y$  is no greater than about 100,000 grams/mole.
- 61. The polymer of claim 57 wherein R<sup>1</sup> and R<sup>2</sup> are each independently a straight chain alkylene group, an arylene group, or combinations thereof.
- 62. The polymer of claim 61 wherein R<sup>1</sup> and R<sup>2</sup> are each independently groups containing up to 100 carbon atoms.
- 63. The polymer of claim 62 wherein R<sup>1</sup> and R<sup>2</sup> are each independently groups containing up to 20 carbon atoms.

Page 12 of 13

Applicant(s): BENZ et al. Serial No.: 10/663.926 Confirmation No.: 2299 Filed: September 16, 2003

For: COMPOUNDS CONTAINING QUATERNARY CARBONS AND SILICON-CONTAINING GROUPS.

MEDICAL DEVICES, AND METHODS

- 64. The polymer of claim 63 wherein R<sup>1</sup> and R<sup>2</sup> are each independently groups containing 2 to 20 carbon atoms.
- 65. The polymer of claim 57 wherein each R2 includes at least two carbon atoms.
- 66. The polymer of claim 57 wherein each R<sup>3</sup> is independently a straight chain alkyl group, an aryl group, or combinations thereof, optionally including heteroatoms.
- 67. The polymer of claim 66 wherein each R<sup>3</sup> is independently a straight chain alkyl group containing 1 to 20 carbon atoms.
- 68. The polymer of claim 57 wherein each Y is OH.
- 69. The polymer of claim 57 wherein each R<sup>4</sup> is independently H or a straight chain alkyl group.
- 70. The polymer of claim 57 which is linear, branched, or crosslinked.
- 76. A method of making a polymer comprising a group of the formula

$$-[-(R^1)_n-(-Z-(R^2)_m-)_p-(-Si(R)_2-V_r-)_s-]_q$$

the method comprising combining an organic compound containing two or more groups capable of reacting with hydroxyl or amine groups with a polymeric starting compound of the formula:

$$Y-[-(R^1)_{a}-(-Z-(R^2)_{m}-)_{p}-(-Si(R)_2-V_{r}-)_{s}-]_{q}-R^5-Y$$

wherein:

each Y is independently OH or  $NR^4H$ ; n = 0 or 1;

Page 13 of 13

Applicant(s): BENZ et al. Serial No.: 10/663,926 Confirmation No.: 2299 Filed: September 16, 2003

For COMPOUNDS CONTAINING QUATERNARY CARBONS AND SILICON-CONTAINING GROUPS,

MEDICAL DEVICES, AND METHODS

m = 0 or 1; p = 1-100,000; r = 0-100,000; s = 1-100,000; q = 1-100,000;

R<sup>1</sup>, R<sup>2</sup>, and R<sup>5</sup> are each independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms;

Z is  $-C(R^3)_2$ - wherein each  $R^3$  is independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms, wherein the two  $R^3$  groups within  $-C(R^3)_2$ - can be optionally joined to form a ring;

each R is independently a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof, optionally including heteroatoms;

each R<sup>4</sup> is independently H or a saturated or unsaturated aliphatic group, an aromatic group, or combinations thereof; and

V is -O-Si(R), or  $R^1$ .